

## **COMPARISON BETWEEN DIRECT AGGLUTINATION TEST AND INDIRECT FLUORESCENT ANTIBODY TEST FOR THE DETECTION OF *Neospora caninum* ANTIBODIES IN NATURALLY EXPOSED DOGS**

WILLIAN A. CANÓN FRANCO<sup>1</sup>, DENISE P. BERGAMASCHI<sup>2</sup>, LAÍS M.A. CAMARGO<sup>3</sup>, VÂNIA S.O. PAULA<sup>1</sup>, SILVIO L.P. SOUZA<sup>1</sup>, SOLANGE M. GENNARI<sup>1\*</sup>

**ABSTRACT.** CANÓN FRANCO, W. A., BERGAMASCHI, D. P., CAMARGO, L. M. A., PAULA, V. S. O., SOUZA, S.L.P., GENNARI, S.M. **Comparison of direct agglutination test and indirect fluorescent antibody test for the detection of *Neospora caninum* antibodies in naturally exposed dogs.** [Comparação entre a aglutinação direta e a imunoflorescência indireta para detecção de anticorpos de *Neospora caninum* em cães naturalmente infectados.] *Revista Brasileira de Parasitologia Veterinária*, v. 12, n. 1, p. 4-6, 2003. Depto de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, Av. Prof. Orlando Marques de Paiva 87, Cidade Universitária, 05508-000, Brazil. E-mail: sgennari@u\_W.br

The indirect fluorescent antibody test (IFAT) and *Neospora agglutination* test (NAT) are two serologic test that detect antibodies to whole tachyzoites. In this study IFAT and NAT were, compared for the detection of *N. caninum* antibodies in naturailly exposed dogs. Sera from 157 dogs from the county of Monte Negro, Rondônia, with 8.3% seropositivity for *N. caninum* using IFAT ( $\geq 50$ ) were analyzed by NAT ( $\geq 25$ ), presenting sensitivity of 61.5% (Confidence Interval 95%: 53.9 to 69.2%), specificity of 89.6% (C.I. 95%: 84.8 to 94.4%) and positiveness of 14.7% (23/157).

**KEY WORDS:** Indirect fluorescent-antibody test, *Neospora agglutination* test, specificity, sensitivity *Neospora caninum*, dogs.

### **RESUMO**

A reação de imunoflorescência indireta (RIFI) e o teste de aglutinação modificada para *Neospora caninun* (NAT) são duas provas utilizadas na detecção de anticorpos anti-*N. caninum*. Neste estudo RIFI e NAT foram comparadas na detecção de anticorpos anti-*N. caninum* em cães naturalmente infectados. Soro de 157 cães do município de Monte Negro, Rondônia, com 8,3% de soropositividade para *N. caninum* pela RIFI ( $\geq 50$ ) foram analisados pelo NAT ( $\geq 25$ ), que apresentou uma sensibilidade de 61,5% (Intervalo de Confiança 95%: 53,9 a 69,2%) e uma especificidade de 89,6% (I.C. 95%: 84,8% a 94,4%) e positividade de 14,7% (23/157).

<sup>1</sup>Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia - Universidade de São Paulo (USP). Av. Prof. Orlando Marques de Paiva, 87, CEP 05508-000 - Cidade Universitária, São Paulo, SP, Brasil. E-mail: sgennari@\_u-W.br

<sup>2</sup>Departamento de Epidemiologia, Faculdade de Saúde Pública, USP.

<sup>3</sup>Departamento de Parasitologia, Instituto de Ciências Biomédicas, USP.

\* Author for correspondence: S. M. Gennari- E-mail: sgennari@\_u-W.br phone: 11-309176-54, fax: 11 3091-7928.

**PALAVRAS-CHAVE:** Reação de Imunoflorescência Indireta, Teste de Aglutinação para *Neospora caninum* especificidade, sensibilidade, *Neospora caninum*, cães.

### **INTRODUCTION**

Antibodies to *Neospora caninum* have been reported in dogs worldwide (DUBEY; LINDSAY, 1996; LINDSAY; DUBEY, 2000); including from Brazil, both in rural and urban areas (SOUZA et al., 2002; GENNARI et al., 2002). A titer of 1:50 is considered indicative of *N. caninum* infection and indirect fluorescent antibody test, (IFAT) has been used as goldstandard to detect *N. caninum* antibodies (DUBEY et al., 1988; BJÖRKMAN; UGGLA, 1999) however, IFAT requires a species specific conjugate and special equipment. *Neospora agglutination* test (NAT) is simple to perform and does not. require species specific conjugate (ROMAND et al., 1998). However, the sensitivity and specificity of this test, in dogs, have not been clearly described. The objective of the present study was to compare the performance of the NAT and IFAT for detecting *N. caninum* antibodies in dogs.

## MATERIAL AND METHODS

Sera from 157 naturally exposed dogs from the county of Monte Negro, state of Rondônia - Brazil, were first tested by the IFAT using culture-derived tachyzoites, of NC-1 isolate (DUBEY et al., 1988b) and the rabbit anti-canine IgG conjugate (Sigma, St. Louis, MO). Sera were tested at 2-fold dilutions starting at 1:50, using the procedure described by Dubey et al. (1988b). The same sera were later evaluated by NAT, performed as described by ROMAND et al. (1998) using formalin fixed whole tachyzoites and tested at 2-fold dilution starting at 1:25 as recommended by Bjorkman and Uggla (1999).

Performance of the NAT was assessed by measures of sensitivity, specificity and global concordance and their respective confidence intervals (C.I.) of 95% (GARDNER; GREINER, 2000) using IFAT as gold test.

## RESULTS

By IFAT antibodies to *N. caninum* were found in 13 (8.3%) of dogs and prevalence study was published elsewhere (CAÑON-FRANCO et al., 2003). Using NAT the seropositivity for antibodies anti-*N. caninum* was 14.7%, with 23 positive dogs. Among the positives ones, 18 (78.30%) dogs presented titer of 25 and 5 (21.7%) titer of 50.

When compared with IFAT ( $\geq 50$ ), the performance study for the NAT ( $\geq 25$ ), to detect *N. caninum* antibodies, showed 61.5% sensitivity (CI 95%: 53.9 to 69.2%) and 89.6% specificity (CI 95%: 84.8 to 94.40%). Global concordance was 87.3%. Fifteen samples that were negative by IFAT were positive by NAT and five samples negatives by NAT were positive by IFAT (Table 1). A fall in sensitivity and a rise in specificity of the test were observed with the increase of the cut-off point as illustrated in Table 2.

Table 1. Detection of *Neospora caninum* antibodies by *Neospora* agglutination test (NAT  $\geq 225$ ) and by indirect fluorescent antibody test (IFAT  $\geq 50$ ), in dogs naturally infected, from the county of Monte Negro, Rondônia.

Tests	IFAT		
	+	-	Total
NAT			
+	8	15	23
-	5	129	134
Total	13	144	157

Table 2. Values for sensitivity and specificity for *Neospora* agglutination test (NAT) compared to indirect fluorescent antibody test (IFAT  $\geq 50$ ) by different NAT cut-off points to detect *Neospora caninum* antibodies.

NAT cut-off point	Sensitivity (%)	Specificity (%)
$\geq 25$	61.5	89.6
$\geq 50$	15.4	97.9

## DISCUSSION

In the present study the NAT performed poorly, owing to low sensitivity (61.5%) for cut-off point  $\geq 25$ . Several cut-off points have been evaluated for the NAT; the most widely used in dogs is  $\geq 25$  (BJÖRKMAN; UGGLA, 1999).

The differences between the performances of NAT versus IFAT may be due to types of antibodies measured, and different epitopes detected. The NAT detects only IgG because the 2-mercaptoethanol used in the test destroys specific and non-specific IgM, whereas IFAT detects both types of antibodies IgG and IgM.

Further studies are needed to compare these tests in experimentally infected dogs.

**Acknowledgments.** The authors gratefully acknowledge the collaboration of colleagues from the Faculdade de Medicina Veterinária e Zootecnia and Instituto de Ciências Biomédicas - USP, who collected the blood samples. We thanks Dr. S. Romand and O. Thulliez for the gift of NAT antigen, and Dr. J. P. Dubey for the comments in the manuscript.

## REFERENCES

- BARKMAN, C., UGGLA, A. Serological diagnosis of *Neospora caninum* infection. *International Journal for Parasitology*, v. 29, p. 1497-1507, 1999.
- CAÑON-FRANCO, W. A., BERGAMASCTU, D. P., LABRU-NA, M. B., CAMARGO, L. M. A., SOUZA, S. L. P., SILVA, J. C. R., PINTER, A., DUBEY, J. P., GENNARI, S. M. Prevalence of antibodies to *Neospora caninum* in dogs from Amazon, Brazil. *Veterinary Parasitology*, 2003. (no prelo).
- DUBEY, J. P., LINDSAY, D. S. A review of *Neospora caninum* and neosporosis. *Veterinary Parasitology*, v. 67, p. 1-59, 1996.
- DUBEY, J. P., CARPENTER, J. L., SPEER C. A., TOPPER, M. J., UGGLA, A. Newly recognized fatal protozoan disease of dogs. *Journal of the American Veterinary Medical Association*, v. 192, p. 1269-1285, 1988.
- DUBEY, J. P., HATTEL, A. L., LINDSAY, D. S., TOPPER, M. J. Neonatal *Neospora caninum* infection in dogs: isolation of the causative agent and experimental transmission. *American Veterinary Medical Association*, v. 193, p. 1259-1263, 1988b.
- GARDNER, I. A., GREINER, M. *Advanced methods for test validation and interpretation in veterinary medicine*, Berlin: Universität Berlin and the University of California Davis, 2000, 102p.
- GENNARI, S. M., YAI, L. E. O., D'ÁURIA, S. N. R., CARDOSO, S. M. S., KWOK, O. C. H., JENKINS, M. C., DUBEY, J. P. Occurrence of *Neospora caninum* antibodies in sera from dogs of the city of São Paulo, Brazil. *Veterinary Parasitology*, v. 106, p. 177-179, 2002.

LINDSAY, D. S., DUBEY, J. P. Canine neosporosis. *Journal of Veterinary Parasitology*, v. 14, p. 1-11, 2000.

ROMAND, S., THULLIEZ, P., DUBEY, J. P. Direct agglutination test for serologic diagnosis of *Neospora caninum* infection. *Parasitology Research*, v. 84, p. 50-53, 1998.

SOUZA, S. L. P., GUIMARÃES, J. S., FERREIRA, F., DUBEY, J. P., GENNARI, S. M. Prevalence of *Neospora caninum* antibodies in dogs from dairy cattle farms in Paraná, Brazil. *Journal of Parasitology*, v. 88, p. 408-409, 2002.

Received on december 17, 2002.

Accepted for publication on August 2, 2003.