

Parasites of economically important bivalves from the southern coast of Bahia State, Brazil

Parasitos de bivalves de interesse econômico no Litoral Sul do Estado da Bahia, Brasil

Gabriela Calvi Zeidan¹; Mariane dos Santos Aguiar Luz²; Guisla Boehs^{1,2*}

¹Programa de Pós-graduação em Sistemas Aquáticos Tropicais, Universidade Estadual de Santa Cruz – UESC, Ilhéus, BA, Brasil

²Programa de Pós-graduação em Ciência Animal, Universidade Estadual de Santa Cruz – UESC, Ilhéus, BA, Brasil

Received June 23, 2012

Accepted October 16, 2012

Abstract

This study investigated the parasites of three commercially important bivalve species (*Crassostrea rhizophorae*, *Mytella guyanensis* and *Lucina pectinata*) from the southern coast of Bahia, Brazil. A total of 540 specimens were collected in August 2009 and February 2010, at three localities. The bivalve specimens were measured on their longest axis, opened, and macroscopically examined for the presence of parasites or signs of disease. They were then fixed in Davidson's solution and subjected to routine histological processing, with paraffin embedding and H&E staining; next, the specimens were examined under a light microscope. No parasites were observed associated with *L. pectinata*. *Rickettsia*-like organisms (RLOs), *Sphenophrya* sp. (Ciliophora), *Nematopsis* sp. (Apicomplexa), *Urastoma* sp. (Turbellaria) and *Bucephalus* sp. (Digenea) were observed in both *C. rhizophorae* and *M. guyanensis*, as well as *Ancistrocoma* sp. (Ciliophora) and *Tylocephalum* sp. (Cestoda) in the former. A high prevalence of *Nematopsis* sp. was seen, but caused no apparent damage to the host. *Bucephalus* sp. caused the destruction of tissues, with castration, but showed low prevalence. The other parasites occurred in low prevalence and intensity, without causing significant damage.

Keywords: *Crassostrea rhizophorae*, *Lucina pectinata*, *Mytella guyanensis*, pathogens, bivalves, parasites.

Resumo

Neste estudo foram investigados os parasitos de três espécies de bivalves de interesse econômico (*Crassostrea rhizophorae*, *Mytella guyanensis* e *Lucina pectinata*) da Bahia. Foram analisados 540 exemplares, obtidos em duas coletas (agosto-2009 e fevereiro-2010), em três localidades. Os bivalves foram medidos quanto ao seu maior eixo, abertos e examinados macroscopicamente quanto à presença de parasitos ou sinais de enfermidades. Depois disso, foram fixados em solução de Davidson e processados por rotina de histologia, com inclusão em parafina e coloração com H&E. O material foi examinado ao microscópio de luz. Nenhum parasito esteve associado a *L. pectinata*. Bactérias do tipo RLOs (organismos assemelhados a *Rickettsia*), *Sphenophrya* sp. (Ciliophora), *Nematopsis* sp. (Apicomplexa), *Urastoma* sp. (Turbellaria) e *Bucephalus* sp. (Digenea) foram observados tanto em *C. rhizophorae* quanto em *M. guyanensis*, sendo que a primeira apresentou ainda *Ancistrocoma* sp. (Ciliophora) e *Tylocephalum* sp. (Cestoda). *Nematopsis* sp. ocorreu em alta prevalência, porém, aparentemente, não causou danos aos bivalves. *Bucephalus* sp. causou destruição de tecidos, com castração, mas foi pouco prevalente. Os demais parasitos ocorreram em baixa prevalência e intensidade de infecção e sem causar danos.

Palavras-chave: *Crassostrea rhizophorae*, *Lucina pectinata*, *Mytella guyanensis*, patógenos, bivalves, parasitas.

Introduction

Many organisms can be found associated with bivalve mollusks, including viruses, bacteria, protists, fungi and metazoans (KINNE, 1983). Some of these symbionts can cause damage and sometimes disease in mollusks, thus affecting natural and farmed stocks.

Crassostrea rhizophorae (Guilding, 1828) (Ostreidae), *Mytella guyanensis* (Lamarck, 1819) (Mytilidae) and *Lucina pectinata* (Gmelin, 1791) (Lucinidae), all prominent species in the coastal mangroves of Bahia state, are economically important and are used for both local consumption and commercial sale in the region.

Crassostrea rhizophorae is already being cultivated on a small scale in Baía de Todos os Santos and in some areas of the southern coast of Bahia, as well as in Valença and Baía de Camamu. Despite

*Corresponding author: Guisla Boehs

Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz – UESC, Rod. Ilhéus-Itabuna, Km 16, CEP 45650-900, Ilhéus, BA, Brasil

e-mail: gboehs@uesc.br

the potential for cultivation of the two other species, they are still exploited only as extractive resources.

According to Rios (2009), the natural geographical distribution of *C. rhizophorae* covers the southern Caribbean, Venezuela, Suriname, Brazil and Uruguay; *M. guyanensis* is found from Mexico to Peru and from Venezuela to Brazil; and *L. pectinata* is distributed from North Carolina to Florida, Texas (USA), West Indies, Venezuela, Suriname and most of the coast of Brazil, from Amapá, in the north, to Santa Catarina, in the south. *Crassostrea rhizophorae* is known along the Brazilian coast as “ostra-do-mangue” (mangrove oyster), attached to hard substrates such as rocks and roots of the red mangrove *Rhizophora mangle* L.. The other two species inhabit unconsolidated substrates (sandy-muddy sediments): *M. guyanensis* lives near mangrove roots, where it burrows just below the surface of the sediment; *L. pectinata* burrows more deeply, from 10 to 20 cm.

Studies on parasites and diseases of mollusks are important since they provide the necessary knowledge for crop management and maintenance of natural resources. As reviewed by Boehs et al. (2012), on the Brazilian coast, the main records of pathogens in bivalve mollusks (e.g., *Crassostrea rhizophorae*, *C. gigas*, *Perna perna*, *Mytella guyanensis*, *Anomalocardia brasiliensis*, *Tagelus plebeius*) include bacteria (e.g., *Rickettsia*-like organisms), protozoans (e.g., *Sphenophrya* sp., *Trichodina* sp., *Ancistrocoma* sp., *Nematopsis* sp., *Perkinsus* sp., *Steinhausia mytilovum*), unidentified fungi, and metazoans (e.g., *Urastoma* sp., *Bucephalus margaritae*, *Tylocephalum* sp., *Polydora websteri*, *Pseudomyicola spinosus*). Yet, these studies are still sparse, including in Bahia. The objective of this study was to catalogue the parasites of three economically important bivalve species (*C. rhizophorae*, *M. guyanensis* and *L. pectinata*) from the southern coast of Bahia, to generate knowledge both for managing natural stocks and for aquaculture activities (especially oyster farming) in the region.

Materials and Methods

Sampling was performed at three localities on the southern coast of Bahia: Camamu (13° 57' S and 39° 03' W), Valença (13° 28' S and 39° 05' W) and Canavieiras (15° 44' S and 38° 55' W), in two sampling campaigns, one in August 2009 (winter) and the other in February 2010 (summer). At each locality, 30 specimens of each species were collected, totaling 180 from an overall total of 540 specimens. The collections were made manually, during low spring tide; oysters were collected from the mangrove roots with the aid of a knife. During each sampling, the temperature and salinity of adjacent waters were measured using a standard mercury thermometer and an Atago S/Mill manual refractometer, respectively. The specimens were brought to the Laboratory of Animal Histology (Universidade Estadual de Santa Cruz, UESC)

in buckets, within a maximum period of six hours, and were immediately processed. The mollusks were washed under tap water and measured on their longest axis (anterioposterior = length in *L. pectinata*, and dorsoventral = height in *M. guyanensis* and *C. rhizophorae*). Then, the specimens were opened with a knife and examined for the presence of parasites or anomalies seen with the naked eye; a cross-section of approximately 5 mm was cut with a scalpel and immersed in Davidson's fixative solution (SHAW; BATTLE, 1957) for 24 hours. The fixed tissues were processed using the routine histological technique, which included: dehydration in a graded ethanol series, diaphanization in xylene, paraffin embedding, and microtome sections of about 7 µm in thickness. The sections were stained with Harris hematoxylin and eosin (H&E) (HOWARD et al., 2004), and mounted under coverslips. Histological preparations were analyzed by light microscopy. After the parasites were identified and their site of infection was recorded, their prevalence was calculated as a function of the number of infected hosts divided by the number of hosts examined for a parasite species, according to Bush et al. (1997); the result was expressed as a percentage. All measurements are in micrometers. The Sørensen qualitative similarity index (ODUM, 1988) was used to compare the parasitic similarity between hosts and localities. In the case of one of the parasites, which occurred in high frequency in the samples, we used the Kruskal-Wallis test ($\alpha = 0.05\%$) to compare the prevalence between localities and sample periods. The intensity of infection by parasites occupying large areas of tissue was calculated through the stereological technique proposed by Lowe et al. (1994), using a Weibel graticule. The tissue area occupied by parasites (PT) was evaluated at five locations of the infected animal's body, with subsequent calculation of the mean. Results were expressed in percentages. For parasites that showed low-level infections, a visual count of the number of parasites/histological section was made. Minimum and maximum values are followed, in parentheses, by arithmetic means \pm standard deviation and the number of measured specimens.

Results

Water temperature ranged between 25 and 31.2 °C at the three sampling localities, in the two periods, and the salinity ranged between 11 and 30‰ (Table 1).

There were no macroparasites or macroscopic signs of disease. Microscopic analyses revealed an absence of parasites in *L. pectinata* and the presence of bacteria, protozoa and metazoans in *C. rhizophorae* and *M. guyanensis*. *Ancistrocoma* sp. (Ciliophora: Ancistrocomidae) and *Tylocephalum* sp. (Cestoda: Lecanicephalidea: Tetragonocephalidae) were seen only in *C. rhizophorae*, while an unidentified metazoan was observed only in *M. guyanensis*. The two species also contained RLOs bacteria (*Rickettsia*-like

Table 1. Datas of temperature (T) and salinity (S) of the three study localities in August 2009 and February 2010, from the southern coast of Bahia State, Brazil.

	Valença		Camamu		Canavieiras	
	T	S	T	S	T	S
August 2009	25.5	11	27.5	29	25	15
February 2010	30.1	29	31.2	29	30	30

organisms), *Sphenophrya* sp. (Ciliophora: Sphenophryidae), *Nematopsis* sp. (Apicomplexa: Eugregarinida: Porosporidae), *Urastoma* sp. (Turbellaria: Urastomidae) and *Bucephalus* sp. (Digenea: Bucephalidae), resulting in a parasitic similarity of 77% between these bivalves. With the exception of *Tylocephalum* sp., which occurred only at Canavieiras, and an unidentified metazoan, occurring exclusively at Valença, the other parasites were recorded at all three localities; this resulted in 86% parasitic similarity between Valença and Canavieiras, and 92% between these localities and Camamu. The parasitic similarity between sampling periods was 86%.

Colonies of intracellular RLOs occurred in the epithelium of the digestive gland of *C. rhizophorae* and *M. guyanensis* (Figure 1a, b), the latter showing a prevalence of up to 30% at Canavieiras (Table 2). The diameter of colonies ranged from 6.6 to 20 μm ($10.4 \pm 4.5 \mu\text{m}$; $n = 10$) in *C. rhizophorae* and 9.5 to 21 μm ($4.2 \pm 5.3 \mu\text{m}$; $n = 4$) in *M. guyanensis*. The observed intensity of infection was 1.2 colonies/histological section in *C. rhizophorae* and 1-6 colonies/histological section in *M. guyanensis*. The RLOs led to localized hypertrophy and cell lysis.

The ciliate *Sphenophrya* sp. occurred in the gill epithelium of *C. rhizophorae* and *M. guyanensis*. This protist measured between 2 and 6 μm in length in *C. rhizophorae* ($3.7 \pm 1.2 \mu\text{m}$; $n = 17$) and between 1 and 6 μm in *M. guyanensis* ($2.6 \pm 1.4 \mu\text{m}$; $n = 17$); in both hosts, the ciliate had an oval shape and a basophilic nucleus. The parasite had a higher prevalence in *C. rhizophorae* (Table 2), where it induced xenoma formation (lesion arising from the proliferation of this protozoa, characterized by prominent hypertrophy of the host cell and its nucleus) in approximately 70% of cases. Xenomas were only recorded in collections performed in August, corresponding to late winter in the region. Xenomas were seen in both hosts an early stage of formation (Figure 1c), containing 2-3 ciliates, and an advanced stage, with about 70-80 ciliates/host cell (Figure 1d); however, the infection intensity was always very low: 1-2 xenomas/histological section. Apparently, damage was restricted to the infected cell.

The ciliate *Ancistrocoma* sp. was only observed in *C. rhizophorae*, in the lumen of its digestive tubules (Figure 1e), with prevalences up to 16.6% (Table 2). This oval-shaped protozoan had a mean length between 5 and 11 μm ($8.6 \pm 1.6 \mu\text{m}$; $n = 7$), with a granular and basophilic macronucleus. The intensity of infection ranged from 1 to 6 ciliates/histological section.

Gregarine oocysts of *Nematopsis* were observed to infect hemocytes (Figure 1f) in both host species *C. rhizophorae* and *M. guyanensis* (prevalence was significantly higher in the former). There were no significant differences in the prevalence of parasites as a function of place and time. Oocysts in *C. rhizophorae* measured between 5.5 and 11 μm ($7.1 \pm 1.4 \mu\text{m}$; $n = 20$) in length, and between 3 and 7 μm (4.9 ± 1.1 ; $n = 20$) in *M. guyanensis*. These parasites occurred mostly in the gills and mantle, and less often in the digestive gland of both hosts. In *C. rhizophorae*, they were also found in the adductor muscle, and in *M. guyanensis* in the foot. The number of oocysts/phagocyte was generally 1-3; however, in *C. rhizophorae*, there were up to 9 oocysts per phagocyte, and in *M. guyanensis*, up to 8 oocysts per phagocyte. Despite their conspicuous prevalence, especially in *C. rhizophorae*, the intensity of infection of these protists was moderate. The relative area

occupied in the host tissue of *C. rhizophorae* was on average 9%, and 7% in *M. guyanensis*. Tissue damage and hemocyte response were not observed.

The turbellarian *Urastoma* sp. was recorded between gill filaments (Figure 2a) of the two host species and also in the mantle cavity of *C. rhizophorae*. In this species, the turbellarian was found at all three sampling localities, with prevalence rates of up to 56.6% in August at Canavieiras (Table 2). The prevalence in *M. guyanensis* was lower (3.3%), and this turbellarian was absent in mollusks from Valença (Table 2). The observed intensities for *C. rhizophorae* and *M. guyanensis* were 1-5, and 1-2 turbellarians/histological section, respectively. This ectocommensal caused no apparent damage.

There was a low prevalence of *Bucephalus* sp. (Figure 2b) for *C. rhizophorae* and *M. guyanensis* at all three localities (Table 2). The parasite was observed at the sporocyst larval stage, when structures (sporocysts) consisted of germinative masses and cercariae. The trematode occurred in the mantle/gonad, digestive gland and gills. The method used to quantify the area occupied by the parasite (PT = Parasite Tissue) in the host tissue (stereology) ranged from 32.3% to 39% in *C. rhizophorae* and 30.4% to 48.5% in *M. guyanensis*, which is considered a high level of infection according to Lowe et al. (1994) (classification stage III = 25-50% PT). One specimen of *M. guyanensis* showed 52.3% PT, indicating very high infection intensity. There were no hemocyte responses; however, all infected specimens showed parasitic castration, with complete destruction of gametes and follicles and impossibility of sex determination; a single specimen of *M. guyanensis* showed a few follicles with vestiges of male gametes in some areas of the gonad.

A metacestode of *Tylocephalum* sp. was identified between the digestive-gland acini of a *C. rhizophorae* specimen from Canavieiras (Table 2). Around the metacestode, hemocyte infiltration was seen, clearly evidencing parasite-induced resorption (Figure 2c). There were two parasites/histological section.

A specimen of *M. guyanensis* collected in August at Valença showed an unidentified metazoan, perhaps a metacercaria, with only one parasite specimen per histological section in the gill filament. Hemocyte infiltration was observed around the parasite, but no tissue damage was apparent (Figure 2d; Table 2).

Discussion

The virtual absence of parasites in *L. pectinata* might be related to its location in the substrate, since this bivalve burrows approximately 20 cm below the surface of the sandy-muddy sediment of mangroves. According to Rios (2009), the conspicuous foot of bivalves of the family Lucinidae, which are tunnel builders, allows them to live in the deeper layers of the substrate (containing hydrogen sulfide and little or no oxygen), where other bivalve species can barely survive. Perhaps, these conditions may limit the establishment of parasites in *L. pectinata*. However, at least the digenean *Bucephalus* sp. has been observed in this species in another locality of the southern Bahia, in low prevalence (1.5%) (OLIVEIRA, 2008).

The high similarity in parasitic fauna between *C. rhizophorae* and *M. guyanensis*, as well as that observed among different localities and




Figure 1. Parasites in two bivalves of economic interest from the south Bahia, Brazil. A-B. Intracellular colonies of *Rickettsia*-like organisms (RLOs) in the epithelium of digestive glands of a) *Mytella guyanensis* and b) *Crassostrea rhizophorae*. Bar = 20 μ m; c-d) *Sphenophrya* sp. in the gills of *Crassostrea rhizophorae* with xenoma formation, being c) in earlier and d) more advanced stage. Arrows = ciliates; * = nucleus of the host cell. Bar = 20 μ m; e) *Ancistrocoma* sp. in the lumen of digestive tubule of *Crassostrea rhizophorae*. Bar = 10 μ m; f) Oocysts of the gregarine *Nematopsis* sp. in connective tissue of the gills of *Crassostrea rhizophorae*. Bar = 20 μ m.

Table 2. Parasites and their prevalences (%) in two bivalves of economic interest of the coastal wetlands from the southern coast of Bahia State, Brazil collected in August 2009 (A) and in February 2010 (F) (n = 60/local).

Parasites	<i>Crassostrea rhizophorae</i>						<i>Mytella guyanensis</i>					
	(A = 5.91 cm DP ± 1.23; n = 180)						(A = 4.78 cm DP ± 0.82; n = 180)					
	Valença		Camamu		Canavieiras		Valença		Camamu		Canavieiras	
	A	F	A	F	A	F	A	F	A	F	A	F
RLOs	6.6	3.3	3.3							6.6	9.9	30
<i>Sphenophrya</i> sp.*	3.3			9.9				3.3	6.6			3.3
<i>Sphenophrya</i> sp.**	20		6.6		3.3							
<i>Ancistrocoma</i> sp.	16.6		6.6			3.3						
<i>Nematopsis</i> sp.	100	93.3	96.6	100	86.6	96.6	76.6	80	53.3	36.6	13.3	26.6
<i>Urastoma</i> sp.		33.3	9.9		56.6				3.3			3.3
<i>Bucephalus</i> sp.				6.6			3.3					9.9
<i>Tylocephalum</i> sp.					3.3							
Metazoan NI							3.3					

A = average height of specimens; RLOs = *Rickettsia*-like Organisms; *Sphenophrya* sp. *without xenoma formation; *Sphenophrya* sp. **with xenoma formation; NI = not identified.




Figure 2. Metazoan in two bivalves of economic interest from south Bahia, Brazil. a) *Urastoma* sp. in gill of *Crassostrea rhizophorae*. Arrow = ocellus. Bar = 50 µm; b) Trematode sporocysts of *Bucephalus* sp. containing germ balls and cercariae (arrows) in mantle of *Crassostrea rhizophorae*. Bar = 20 µm; c) *Tylocephalum* sp. metacystode (arrows) in the connective tissue of the digestive gland of *Crassostrea rhizophorae*. * = myzorhynchus. Bar = 50 µm; d) Unidentified metazoan in the gills of *Mytella guyanensis*. Bar = 20 µm.

sample periods, were clearly shown in this study. With respect to the RLOs found in both species, the site of infection in the host (the digestive gland), the localized effect, and the absence of a hemocyte response converge with results from recent studies conducted by Da Silva et al. (2011) and Sabry et al. (2011) on *C. rhizophorae* and *C. gigas* (Thunberg, 1793) in Santa Catarina state, and by Boehs et al. (2010) and Ceuta and Boehs (2012) on *M. guyanensis* in Bahia state. Except for the study by Sabry et al. (2011), who identified prevalences of up to 30% for *C. gigas* (coinciding with the results of this study for *M. guyanensis* collected at Canavieiras, in February 2010), other studies have indicated a low prevalence of these bacteria. Other studies have also found a significant lack of damage, as seen in this study. Yet, Sabry et al. (2011), who observed RLOs in the stomach of one *C. rhizophorae* specimen, noted gastric-epithelium changes caused by these organisms.

Ciliates are commonly found in bivalves, but do not normally cause major damage to their hosts (BOWER et al., 1994). Regarding *Ancistrocoma* sp., the low infection intensity and the apparent lack of damage to the host are in line with previous observations of Sabry et al. (2011) on *C. rhizophorae* and *C. gigas* from Santa Catarina, but with prevalences of up to 40%. As for *Sphenophrya*, according to Bower et al. (1994), these ciliates are found in a variety of bivalve species, in which they may cause xenoma. This lesion was previously observed by Boehs et al. (2009) for *C. rhizophorae* at Baía de Camamu. Other studies of Brazilian coastal mollusks have also reported the occurrence of *Sphenophrya* sp. in oysters, but without xenoma formation (NASCIMENTO et al., 1986; SABRY et al., 2011). In turn, in *C. virginica* (Gmelin, 1791), Winstead et al. (2004) observed xenoma formation in all oysters containing this protist. As observed in this study, the prevalence levels seen by these authors were low; still, the study conducted by Sabry et al. (2011) showed a prevalence of up to 70% for *C. gigas*, at one locality in Santa Catarina.

The high prevalence of the gregarine *Nematopsis*, especially in *C. rhizophorae*, is in line with several studies involving coastal bivalves from Brazil, such as on *C. rhizophorae* (SABRY et al., 2007, 2011; DA SILVA et al., 2011), *M. guyanensis* (PINTO; BOEHS, 2008; BOEHS et al., 2010; CEUTA; BOEHS, 2012), *Anomalocardia brasiliiana* (Gmelin, 1791) (BOEHS et al., 2010) and *Perna perna* (Linnaeus, 1758) (DA SILVA et al., 2011). This study also confirmed that gills and mantle were the main organs affected by this parasite, although it may also occur in other anatomical regions such as the digestive gland and foot. Ceuta and Boehs (2012) also noted a high level of infection by *Nematopsis* sp. in the labial palps, and structural changes in these organs. This parasite is generally reported to cause little damage to bivalves, especially when the infection level is low (BOWER et al., 1994).

Among the observed platyhelminths, the turbellarian *Urastoma* has been reported to live in the gills, mantle cavity and digestive gland of several bivalves (LAUCKNER, 1983). In this study, *Urastoma* was found between the gill filaments of *M. guyanensis* and *C. rhizophorae*, as well as in the mantle cavity of the latter, where prevalence levels reached up to 56.6%. In a study in southern Brazil, Sabry et al. (2011) reported a low prevalence of this turbellarian in the gill region of *C. rhizophorae* and *C. gigas*. Investigating the presence of *U. cyprinae* (Graff, 1882) in *Mytilus galloprovincialis* (Lamarck, 1819), in Galicia (Spain), Robledo et al. (1994)

noted macroscopic lesions in the form of white spots. Through microscopic analysis, these authors also identified disorganization and compression of the gill filaments, hypertrophy of epithelial cells, and hemocyte infiltration, concluding that this ectocommensal causes significant changes in the feeding function. In southern Portugal, Francisco et al. (2010) also observed disorganization of the gill tissues and labial palps of the same host. The lack of apparent damage in this study was probably due to the low infestation intensity in both species.

Similarly to other Bucephalidae, the digenean *Bucephalus* spp. has a complex life cycle, using bivalve mollusks as intermediate hosts and some bony fishes as definitive hosts (LAUCKNER, 1983). These trematodes are so named due to their forked tail with a very short and wide base during the cercaria phase (WARDLE, 1990). Studies in some species of Mytilidae (LASIAK, 1993; BOEHS et al., 2010; CEUTA; BOEHS, 2012) have found macroscopic signs such as an orange color and/or fibrous appearance of the mantle, as indications of the presence of sporocysts, which could not be confirmed in this study. The prevalence levels usually below 10% corroborated the results from several studies involving Mytilidae on the Brazilian coast (DA SILVA et al., 2002, 2011; BOEHS et al., 2010;), the oyster *C. rhizophorae* from the Brazilian coast (NASCIMENTO et al., 1986), *C. virginica* from Florida, USA (WINSTEAD et al., 2004) and *M. galloprovincialis* from Portugal (FRANCISCO et al., 2010). All these studies confirmed tissue damage by Bucephalidae, especially in the gonads, characterizing parasitic castration. Castration was also caused by trematode sporocysts in *Cerastoderma edule* (Linnaeus, 1758) (Cardiidae) (CARBALLAL et al., 2001) and *M. galloprovincialis* (VILLALBA et al., 1997) in the region of Galicia. In this study, the observed infection level and lesions on the digestive gland and gills were smaller than those of the gonad. Despite the damage caused by infection, there was no apparent inflammatory response in the host. Contrary to what was seen in this study, Boehs et al. (2010) observed hemocyte infiltration around dead or degenerating sporocysts in *M. guyanensis*, while Da Silva et al. (2002) reported high hemocyte infiltration in heavily infected *Perna perna*. According to Bower (1992) and Carballal et al. (2001), high infection levels by sporocysts can reduce recruitment at the population level. To date, despite the obvious risk of *Bucephalus* spp. to its hosts, the low prevalence of this parasite does not appear to affect the natural populations of these bivalves on the southern coast of Bahia.

In Brazil, the metacestode *Tylocephalum* sp. has been reported to use different species of bivalves as an intermediate host (NASCIMENTO et al., 1986; BOEHS; MAGALHÃES, 2004; SABRY; MAGALHÃES, 2005; SABRY et al., 2007; BOEHS et al., 2010; DA SILVA et al., 2011; CEUTA; BOEHS, 2012). According to Lauckner (1983), this cestode completes its life cycle in elasmobranchs, and is more common in tropical and subtropical regions. Previous studies involving *C. rhizophorae* indicate the digestive gland as the major site of occurrence of this parasite (NASCIMENTO et al., 1986; SABRY; MAGALHÃES, 2005; SABRY et al., 2007); also, all studies showed encapsulation with intense hemocyte infiltration and in some cases, as in the present study, resorption. The same responses were observed in *Tapes semidecussata* Reeve, 1864 (Veneridae) (CHENG; RIFKIN, 1968) in Hawaii, and in *C. virginica* (WINSTEAD et al., 2004) in Florida.

Conclusion

This study showed that, up to the present, natural populations of *C. rhizophorae* and *M. guyanensis* from the southern coast of Bahia are not seriously affected by the presence of parasites, since most of them were found in low prevalence; also, despite the high prevalence of *Nematopsis* sp., the intensity of infection was low. In spite of the obvious damage that *Bucephalus* sp. causes to tissues, the prevalence of this species was low.

Acknowledgements

The authors are thankful to the National Council for Scientific and Technological Development – CNPq (MCT-Brazil; ID 477317/2008-1) for financially aiding the project, to FAPESB, for granting a master's degree fellowship to the first author, and to CNPq, for granting a scholarship for scientific initiation to the second author.

References

- Boehs G, Lenz TM, Villalba A. Xenomas in *Crassostrea rhizophorae* (Ostreidae) from Camamu Bay, Bahia, Brazil. *Braz J Biol* 2009; 69(2): 457-458. PMID:19675953. <http://dx.doi.org/10.1590/S1519-69842009000200032>
- Boehs G, Magalhães ARM. Simbiontes associados com *Anomalocardia brasiliensis* (Gmelin) (Mollusca, Bivalvia, Veneridae) na Ilha de Santa Catarina e região continental adjacente, Santa Catarina, Brasil. *Rev Bras Zool* 2004; 21(4): 865-869. <http://dx.doi.org/10.1590/S0101-81752004000400021>
- Boehs G, Magalhães ARM, Sabry RC, Ceuta LO. Parasitos e patologias de bivalves marinhos de importância econômica da costa brasileira. In: Silva-Souza AT, Lizama MLA, Takemoto R, organizadores. *Patologia e Sanidade de Organismos Aquáticos*. Maringá: ABRAPOA; 2012, p. 165-194.
- Boehs G, Villalba A, Ceuta LO, Luz JR. Parasites of three commercially exploited bivalve mollusc species of the estuarine region of the Cachoeira river (Ilhéus, Bahia, Brazil). *J Invertebr Pathol* 2010; 103(1): 43-47. PMID:19850046. <http://dx.doi.org/10.1016/j.jip.2009.10.008>
- Bower SM. Diseases and parasites of mussels. In: Gosling E. *Developments in Aquaculture and Fisheries Science: The mussel Mytilus: Ecology, Physiology, Genetics and Culture* (Dev Aquacult Fish Sci, 25). Amsterdam: Elsevier Science Publishers; 1992.
- Bower SM, McCladdery SE, Price IM. Synopsis of infectious diseases and parasites of commercially exploited shellfish. *Annu Rev Fish Dis* 1994; 4(1): 1-199. [http://dx.doi.org/10.1016/0959-8030\(94\)90028-0](http://dx.doi.org/10.1016/0959-8030(94)90028-0)
- Bush AO, Lafferty KD, Lotz JM, Shostak AW. Parasitology meets ecology on its own terms: Margolis et al. revisited. *J Parasitol* 1997; 83(4): 575-583. PMID:9267395. <http://dx.doi.org/10.2307/3284227>
- Carballal MJ, Iglesias D, Santamarina J, Ferro-Soto B, Villalba A. Parasites and pathologic conditions of the cockle *Cerastoderma edule* populations of the Coast of Galicia (NW Spain). *J Invertebr Pathol* 2001; 78(1): 87-97. PMID:11812111. <http://dx.doi.org/10.1006/jipa.2001.5049>
- Ceuta LO, Boehs G. Parasites of mangrove mussel *Mytella guyanensis* (Bivalvia: Mytilidae) of Camamu Bay, Bahia, Brazil. *Braz J Biol* 2012; 72(3): 421-427. PMID:22990810. <http://dx.doi.org/10.1590/S1519-69842012000300002>
- Cheng TC, Rifkin E. The occurrence and resorption of *Tylocephalum* metacestodes in the clam *Tapes semidecussata*. *J Invertebr Pathol* 1968; 10(1): 65-69. [http://dx.doi.org/10.1016/0022-2011\(68\)90264-4](http://dx.doi.org/10.1016/0022-2011(68)90264-4)
- Da Silva PM, Magalhães ARM, Barracco MA. Pathologies in commercial bivalve species from Santa Catarina State, southern Brazil. *J Mar Biol Assoc UK* 2011; 92(3): 571-579. <http://dx.doi.org/10.1017/S0025315411001007>
- Da Silva PM, Magalhães ARM, Barracco MA. Effects of *Bucephalus* sp. (Trematoda: Bucephalidae) on *Perna perna* mussels from a culture station in Ratonas Grande Island, Brazil. *J Invertebr Pathol* 2002; 79(3): 154-162. [http://dx.doi.org/10.1016/S0022-2011\(02\)00026-5](http://dx.doi.org/10.1016/S0022-2011(02)00026-5)
- Francisco CJ, Hermida MA, Santos MJ. Parasites and symbionts from *Mytilus galloprovincialis* (Lamarck, 1819) (Bivalves: Mytilidae) of the Aveiro Estuary Portugal. *J Parasitol* 2010; 96(1): 200-205. PMID:19785477. <http://dx.doi.org/10.1645/GE-2064.1>
- Howard DW, Lewis EJ, Keller BJ, Smith CS. Histological techniques for marine bivalve mollusks and crustaceans. *NOAA Technical Memorandum NCCOS* 2004; 5(1): 1-218.
- Kinne O. *Diseases of Marine Animals*. Hamburg: Biologische Anstalt Helgoland; 1983.
- Lasiak TA. Bucephalid trematode infections in the brown mussel *Perna perna* (Bivalvia: Mytilidae). *Afr J Mar Sci* 1993; 13(1): 127-134. 1993.
- Lauckner G. Diseases of Mollusca: Bivalvia. In: Kinne O. *Diseases of Marine Animals*. Hamburg: Biologische Anstalt Helgoland; 1983. p. 477-970.
- Lowe DM, Salkeld PN, Carr MR. The effect of geographical location on the cellular composition of the mantle tissue of the mussel, *Mytilus edulis*. *J Mar Biol Assoc UK* 1994; 74(1): 225-232. <http://dx.doi.org/10.1017/S0025315400035785>
- Nascimento IA, Smith DH, Kern II F, Pereira SA. Pathological findings in *Crassostrea rhizophorae* from Todos os Santos Bay, Bahia, Brazil. *J Invertebr Pathol* 1986; 47(3): 340-349. [http://dx.doi.org/10.1016/0022-2011\(86\)90105-9](http://dx.doi.org/10.1016/0022-2011(86)90105-9)
- Odum EP. *Ecologia*. Rio de Janeiro: Guanabara Koogan; 1988.
- Oliveira, JB. *Parasitos associados à lambreta Lucina pectinata* (Gmelin, 1791) (Mollusca: Bivalvia) na região estuarina do Rio Cachoeira (Ilhéus, Bahia) [Monografia]. Ilhéus: Universidade Estadual de Santa Cruz; 2008. Disponível na Biblioteca Central da UESC.
- Pinto TR, Boehs G. *Nematopsis* sp. (Apicomplexa: Eugregarinida) em *Mytella guyanensis* (Lamarck, 1819) (Bivalvia: Mytilidae) da região estuarina do Rio Cachoeira, Ilhéus, Bahia, Brasil. *Braz J Vet Res Anim Sci* 2008; 45(2): 95-100.
- Rios E. *Compendium of Brazilian Sea shells*. Rio Grande: Evangraf; 2009.
- Robledo JAF, Cáceres-Martínez J, Sluys R, Figueras A The parasitic turbellarian *Urastoma cyprinae* (Platyhelminthes: Urastomidae) from blue mussel *Mytilus galloprovincialis* in Spain: occurrence and pathology. *Dis Aquat Org* 1994; 18(1): 203-210. <http://dx.doi.org/10.3354/dao018203>
- Sabry RC, Da Silva PM, Gesteira TCV, Pontinha VA, Magalhães ARM. Pathological study of oysters *Crassostrea gigas* from culture and *C. rhizophorae* from natural stock of Santa Catarina Island, SC, Brazil. *Aquaculture* 2011; 320(1-2): 43-50. <http://dx.doi.org/10.1016/j.aquaculture.2011.08.006>

Sabry RC, Gesteira TCV, Boehs G. First record of parasitism in the mangrove oyster *Crassostrea rhizophorae* (Bivalvia: Ostreidae) at Jaguaribe River estuary - Ceará, Brazil. *Braz J Biol* 2007; 67(4): 755-758. PMID:18278331. <http://dx.doi.org/10.1590/S1519-69842007000400024>

Sabry RC, Magalhães ARM. Parasitas em ostras de cultivo (*Crassostrea rhizophorae* e *Crassostrea gigas*) da Ponta do Sambaqui, Florianópolis, SC. *Arq Bras Med Vet Zootec* 2005; 57(S2): 194-203.

Shaw BL, Battle HI. The gross and microscopic anatomy of the digestive tract of the oyster *Crassostrea virginica* (Gmelin). *Can J Zool* 1957; 35(1): 325-347. <http://dx.doi.org/10.1139/z57-026>

Villalba A, Mourelle SG, Carballal MJ, López C. Symbionts and diseases of farmed mussels *Mytilus galloprovincialis* throughout the culture process in the Rías of Galicia (NW Spain). *Dis Aquat Org* 1997; 31(2): 127-139. <http://dx.doi.org/10.3354/dao031127>

Wardle WJ. Larval Bucephalids (Trematoda: Digenea) parasitizing bivalve molluscs in the Galveston Bay Area, Texas. *J Helminthol Soc Wash* 1990; 57(1): 5-11.

Winstead JT, Volety AK, Tolley SG. Parasitic and symbiotic fauna in oysters (*Crassostrea virginica*) collected from the Caloosahatchee River and estuary in Florida. *J Shellfish Res* 2004; 23(3): 831-840.